

# Probability and Random Processes

## ECS 315

Asst. Prof. Dr. Prapun Suksompong

[prapun@siit.tu.ac.th](mailto:prapun@siit.tu.ac.th)

### 1 Probability and You



**Office Hours:**

**BKD 3601-7**

**Monday            14:00-16:00**

**Wednesday       14:40-16:00**

**Everything** we do, everything that happens around us, obeys the laws of **probability**.

We can no more escape them than we can escape gravity... "Probability," a philosopher (Bishop Butler) once said, "is the **very guide of life**."

We are all gamblers who go through life making countless bets on the outcome of countless actions.

**Life is random**

In 2005, this statement (which is true)

**Life is random**

showed up all over the world...

**iPod shuffle**  
**Life is random.**




# Applications of Probability Theory

- The subject of probability can be traced back to the **17th century** when it arose out of the study of **gambling games**.
- The range of applications extends beyond games into business decisions, insurance, law, **medical tests**, and the social sciences.
- The **stock market**, “the largest casino in the world,” cannot do without it.
- The **telephone network**, call centers, and airline companies with their randomly fluctuating loads could not have been economically designed without probability theory.




# “The Perfect Thing”



What is this?



# “The Perfect Thing”



[Share your own customer images](#)

[Search inside this book](#)

Start reading [The Perfect Thing](#) on

## The Perfect Thing: How the iPod Shuffles Commerce, Culture, and Coolness

[Hardcover]

[Steven Levy](#) (Author)

★★★★☆ (27 customer reviews) | Like (1)

Available from [these sellers](#).

**27 new** from \$0.95    **96 used** from \$0.01

Formats	Amazon Price	New from	Used from
Kindle Edition	--	\$10.99	--
Hardcover, Bargain Price	\$10.00	\$10.00	\$3.94
Hardcover, October 24, 2006	--	\$0.95	\$0.01
Paperback	\$14.00	\$0.77	\$0.01
Audible Audio Edition, Abridged	\$17.95	or Free with Audible 30-day free trial	



# What about the shuffle function?

Electronics >  
**About.com** iPhone / iPod  
Part of The New York Times Company

Home iPhone / iPod New to iPhone? Apps iPho

## Is iTunes' Shuffle Mode Truly Random?

By Sam Costello, About.com Guide

<http://ipod.about.com/od/advanceditunesuse/a/itunes-random.htm>



CNET | Australia

Home Mobile Phones Computers Cameras TVs

MP3 Players > News > iTunes: Just how random is random?

## iTunes: Just how random is random?

By David Braue | March 8, 2007 | 127

<http://www.cnet.com.au/itunes-just-how-random-is-random-339274094.htm>

howstuffworks

Adventure Auto Culture Entertainment Home & Garden

Computer | Electronics

## iPod Shuffle Problems: How Random is the iPod Shuffle?

So just how random is the shuffle capability on an iPod Shuffle? Even before the device debuted in 2005, people have wondered about the shuffle function on iPods. Many complain that what they hear from

<http://electronics.howstuffworks.com/ipod-shuffle2.htm>



# USA Currency Coins

- Penny = 1 cent  
(Abraham Lincoln)



- Dime = 10 cents  
(Franklin D. Roosevelt)



- Nickel = 5 cents  
(Thomas Jefferson)



- Quarter = 25 cents  
(George Washington)



# Thai Coins

Front



Back



# randi function


```
>> randi([0,1],1,5)
ans =
     1     1     0     1     1
```

- Generate uniformly distributed **pseudorandom integers**
- `randi(imax)` returns a scalar value between 1 and `imax`.
- `randi(imax,m,n)` and `randi(imax,[m,n])` return an  $m$ -by- $n$  matrix containing pseudorandom integer values drawn from the discrete uniform distribution on the interval  $[1,imax]$ .
  - `randi(imax)` is the same as `randi(imax,1)`.
- `randi([imin,imax],...)` returns an array containing integer values drawn from the discrete uniform distribution on the interval  $[imin,imax]$ .



# Coin Tossing: Relative Frequency

```
close all; clear all;  
N = 1e3; % Number of trials (number of times that the coin is tossed)  
s = randi([0,1],1,N); % Generate a sequence of N Coin Tosses.  
% The results are saved in a row vector s.  
NH = cumsum(s); % Count the number of heads  
plot(NH./(1:N),'LineWidth',1.5); grid on % Plot the relative frequencies
```




# Coin Tossing: Relative Freq. vs. #H-#T

This statement does not say that the difference between #H and #T will be close to 0.

If a fair coin is flipped a large number of times, the **proportion** of heads will tend to get closer to  $1/2$  as the number of tosses increases.






The **difference** between #H and #T will **not** converge to 0.



# Another trial

Relative Freq.

#H-#T





# Another trial

Relative Freq.

#H-#T

