

HW 2 — Due: Sep 2, 9:19 AM (in tutorial session)

Lecturer: Prapun Suksompong, Ph.D.

Instructions

- (a) ONE part of a question will be graded (5 pt). Of course, you do not know which part will be selected; so you should work on all of them.
- (b) It is important that you try to solve all problems. (5 pt)
The extra question at the end is optional.
- (c) Late submission will be heavily penalized.

Problem 1. (Classical Probability and Combinatorics) Shuffle a deck of cards and cut it into three piles. What is the probability that (at least) a court card will turn up on top of one of the piles.

Hint: There are 12 court cards (four jacks, four queens and four kings) in the deck.

Problem 2. (Classical Probability) There are three buttons which are painted red on one side and white on the other. If we tosses the buttons into the air, calculate the probability that all three come up the same color.

Remarks: A *wrong* way of thinking about this problem is to say that there are four ways they can fall. All red showing, all white showing, two reds and a white or two whites and a red. Hence, it seems that out of four possibilities, there are two favorable cases and hence the probability is $1/2$.

Problem 3. Each of the possible five outcomes of a random experiment is equally likely. The sample space is $\{a, b, c, d, e\}$. Let A denote the event $\{a, b\}$, and let B denote the event $\{c, d, e\}$. Determine the following:

- (a) $P(A)$
- (b) $P(B)$
- (c) $P(A^c)$

(d) $P(A \cup B)$

(e) $P(A \cap B)$

[Montgomery and Runger, 2010, Q2-54]

Problem 4. If A , B , and C are disjoint events with $P(A) = 0.2$, $P(B) = 0.3$ and $P(C) = 0.4$, determine the following probabilities:

(a) $P(A \cup B \cup C)$

(b) $P(A \cap B \cap C)$

(c) $P(A \cap B)$

(d) $P((A \cup B) \cap C)$

(e) $P(A^c \cap B^c \cap C^c)$

[Montgomery and Runger, 2010, Q2-75]

Problem 5. The sample space of a random experiment is $\{a, b, c, d, e\}$ with probabilities 0.1, 0.1, 0.2, 0.4, and 0.2, respectively. Let A denote the event $\{a, b, c\}$, and let B denote the event $\{c, d, e\}$. Determine the following:

(a) $P(A)$

(b) $P(B)$

(c) $P(A^c)$

(d) $P(A \cup B)$

(e) $P(A \cap B)$

[Montgomery and Runger, 2010, Q2-55]

Extra Question

Here is an optional question for those who want more practice.

Problem 6. (Combinatorics) Consider the design of a communication system in the United States.

- (a) How many three-digit phone prefixes that are used to represent a particular geographic area (such as an area code) can be created from the digits 0 through 9?
- (b) How many three-digit phone prefixes are possible in which no digit appears more than once in each prefix?
- (c) As in part (a), how many three-digit phone prefixes are possible that do not start with 0 or 1, but contain 0 or 1 as the middle digit?

[Montgomery and Runger, 2010, Q2-45]