

HW 1 — Due: Feb 5

Lecturer: Asst. Prof. Dr. Prapun Suksompong

Instructions

- (a) Must solve all non-optional problems. (5 pt)
 - (i) Write your first name and the last three digit of your student ID on the upper-right corner of *every* submitted page.
 - (ii) For each part, write your explanation/derivation and answer in the space provided.
- (b) ONE part of a question will be graded (5 pt). Of course, you do not know which part will be selected; so you should work on all of them.
- (c) Late submission will be rejected.
- (d) **Write down all the steps** that you have done to obtain your answers. You may not get full credit even when your answer is correct without showing how you get your answer.

Problem 1 (HW1-2015-2). Consider the code $\{0, 01\}$

- (a) Is it nonsingular?

- (b) Is it uniquely decodable?

- (c) Is it prefix-free?

Problem 2 (HW1-2015-2). Consider the random variable X whose support S_X contains seven values:

$$S_X = \{x_1, x_2, \dots, x_7\}.$$

Their corresponding probabilities are given by

x	x_1	x_2	x_3	x_4	x_5	x_6	x_7
$p_X(x)$	0.49	0.26	0.12	0.04	0.04	0.03	0.02

- (a) Find the entropy $H(X)$.
- (b) Find a binary Huffman code for X .
- (c) Find the expected codelength for the encoding in part (b).

Problem 3 (HW1-2015-2). Find the entropy and the binary Huffman code for the random variable X with pmf

$$p_X(x) = \begin{cases} \frac{x}{21}, & x = 1, 2, \dots, 6, \\ 0, & \text{otherwise.} \end{cases}$$

Also calculate $\mathbb{E}[\ell(X)]$ when Huffman code is used.

Problem 4 (HW1-2015-2). These codes cannot be Huffman codes. Why?

- (a) $\{00, 01, 10, 110\}$
- (b) $\{01, 10\}$
- (c) $\{0, 01\}$

Problem 6 (HW1-2015-2). (Optional) The following claim is sometimes found in the literature:

“It can be shown that the length $\ell(x)$ of the Huffman code of a symbol x with probability $p_X(x)$ is always less than or equal to $\lceil -\log_2 p_X(x) \rceil$ ”.

Even though it is correct in many cases, this claim is not true in general.

Find an example where the length $\ell(x)$ of the Huffman code of a symbol x is greater than $\lceil -\log_2 p_X(x) \rceil$.

Hint: Consider a pmf that has the following four probability values $\{0.01, 0.30, 0.34, 0.35\}$.

Problem 7 (HW1-2015-2). (Optional) Construct a random variable X (by specifying its pmf) whose corresponding Huffman code is $\{0, 10, 11\}$.