

EES 315: Probability and Random Processes**2020/1****HW 5 — Due: Not Due***Lecturer: Prapun Suksompong, Ph.D.***Problem 1.**

(a) Suppose that $P(A|B) = 0.4$ and $P(B) = 0.5$ Determine the following:

(i) $P(A \cap B)$

(ii) $P(A^c \cap B)$

[Montgomery and Runger, 2010, Q2-105]

(b) Suppose that $P(A|B) = 0.2$, $P(A|B^c) = 0.3$ and $P(B) = 0.8$ What is $P(A)$? [Montgomery and Runger, 2010, Q2-106]

Problem 4. You have two coins, a fair one with probability of heads $\frac{1}{2}$ and an unfair one with probability of heads $\frac{1}{3}$, but otherwise identical. A coin is selected at random and tossed, falling heads up. How likely is it that it is the fair one? [Capinski and Zastawniak, 2003, Q7.28]

Problem 5. You have three coins in your pocket, two fair ones but the third biased with probability of heads p and tails $1 - p$. One coin selected at random drops to the floor, landing heads up. How likely is it that it is one of the fair coins? [Capinski and Zastawniak, 2003, Q7.29]

Extra Questions

Here are some optional questions for those who want more practice.

Problem 6. Someone has rolled a fair dice twice. Suppose he tells you that “one of the rolls turned up a face value of six”. What is the probability that the other roll turned up a six as well? [Tijms, 2007, Example 8.1, p. 244]

Hint: Note the followings:

- The answer is not $\frac{1}{6}$.
- Although there is no use of the word “given” or “conditioned on” in this question, the probability we seek is a conditional one. We have an extra piece of information because we know that the event “one of the rolls turned up a face value of six” has occurred.

- The question says “one of the rolls” without telling us which roll (the first or the second) it is referring to.

Problem 7.

- (a) Suppose that $P(A|B) = 1/3$ and $P(A|B^c) = 1/4$. Find the range of the possible values for $P(A)$.

- (b) Suppose that C_1, C_2 , and C_3 partition Ω . Furthermore, suppose we know that $P(A|C_1) = 1/3$, $P(A|C_2) = 1/4$ and $P(A|C_3) = 1/5$. Find the range of the possible values for $P(A)$.

Problem 8. In his book *Chances: Risk and Odds in Everyday Life*, James Burke says that there is a 72% chance a polygraph test (lie detector test) will catch a person who is, in

fact, lying. Furthermore, there is approximately a 7% chance that the polygraph will falsely accuse someone of lying. [Brase and Brase, 2011, Q4.2.26]

- (a) If the polygraph indicated that 30% of the questions were answered with lies, what would you estimate for the actual percentage of lies in the answers?

- (b) If the polygraph indicated that 70% of the questions were answered with lies, what would you estimate for the actual percentage of lies?

Problem 9. Software to detect fraud in consumer phone cards tracks the number of metropolitan areas where calls originate each day. It is found that 1% of the legitimate users originate calls from two or more metropolitan areas in a single day. However, 30% of fraudulent users originate calls from two or more metropolitan areas in a single day. The proportion of fraudulent users is 0.01%. If the same user originates calls from two or more metropolitan areas in a single day, what is the probability that the user is fraudulent? [Montgomery and Runger, 2010, Q2-144]

Problem 10. An article in the British Medical Journal [“Comparison of Treatment of Renal Calculi by Operative Surgery, Percutaneous Nephrolithotomy, and Extracorporeal Shock Wave Lithotripsy” (1986, Vol. 82, pp. 879-892)] provided the following discussion of success rates in kidney stone removals. Open surgery (OS) had a success rate of 78% (273/350) while a newer method, percutaneous nephrolithotomy (PN), had a success rate of 83% (289/350). This newer method looked better, but the results changed when stone diameter was considered. For stones with diameters less than two centimeters, 93% (81/87) of cases of open surgery were successful compared with only 87% (234/270) of cases of PN. For stones greater than or equal to two centimeters, the success rates were 73% (192/263) and 69% (55/80) for open surgery and PN, respectively. Open surgery is better for both stone sizes, but less successful in total. In 1951, E. H. Simpson pointed out this apparent contradiction (known as Simpson’s Paradox) but the hazard still persists today. Explain how open surgery can be better for both stone sizes but worse in total. [Montgomery and Runger, 2010, Q2-115]