

Lecture 2

Tuesday, November 03, 2009
10:33 AM

A. Trig. Review

$$\text{Euler: } e^{j\alpha} = \cos \alpha + j \sin \alpha$$

$$\cos \alpha = \frac{1}{2} (e^{j\alpha} + e^{-j\alpha})$$

$$\sin \alpha = \frac{1}{2j} (e^{j\alpha} - e^{-j\alpha})$$

Can use these to derive many formula.

Example:

$$\begin{aligned} \cos A \cdot \cos B &= \frac{1}{2} (e^{jA} + e^{-jA}) \times \frac{1}{2} (e^{jB} + e^{-jB}) \\ &= \frac{1}{4} \left(\underbrace{e^{j(A+B)} + e^{-j(A+B)}}_{2 \cos(A+B)} + \underbrace{e^{j(A-B)} + e^{-j(A-B)}}_{2 \cos(A-B)} \right) \\ &= \frac{1}{2} (\cos(A+B) + \cos(A-B)) \end{aligned}$$

$$A = B$$

$$\cos^2 A = \frac{1}{2} (\cos(2A) + 1)$$

$$\sin^2 A = 1 - \cos^2 A = 1 - \frac{1}{2} \cos 2A - \frac{1}{2} = \frac{1}{2} - \frac{1}{2} \cos 2A$$

B. Fourier Transform: Given $x(t)$

$$X_1(f) = \int_{-\infty}^{\infty} x(t) e^{-j2\pi f t} dt$$

$$X_2(\omega) = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt$$

$$\omega = 2\pi f$$

Q: What is the relationship btw. $X_1(f)$ and $X_2(\omega)$?

$$X_1(f) = X_2(\omega) \Big|_{\omega = 2\pi f} = X_2(2\pi f)$$

$$X_2(\omega) = X_1(f) \Big|_{f = \frac{\omega}{2\pi}} = X_1\left(\frac{\omega}{2\pi}\right)$$

Example Suppose we know that

$$e^{j\omega_0 t} \xrightarrow{\mathcal{F}} 2\pi \delta(\omega - \omega_0)$$

↓

$$e^{j2\pi f_0 t} \xrightarrow{\mathcal{F}} 2\pi \delta(\underline{2\pi f} - \underline{2\pi f_0})$$

$$= \delta(f - f_0)$$

$$\delta(at) = \frac{1}{|a|} \delta(t)$$

Remark: $X(f)$ is usually complex-valued.

Suppose $x(t)$ is real-valued.

$$X(f) = \int x(t) e^{-j2\pi f t} dt$$

$$X(-f) = \int x(t) e^{-j2\pi(-f)t} dt$$

$$= \int x(t) e^{(-1)(j2\pi f t)} dt$$

$$= \int x^*(t) (e^{-j2\pi f t})^* dt$$

$$= \left(\int x(t) e^{-j2\pi f t} dt \right)^*$$

$$= (X(f))^*$$

Conjugate real
valued.
↙ ↘
 $a + jb$

$$= (a + jb)^*$$

$$= a - jb$$

$$x(t) \xrightarrow{\mathcal{F}} X(f)$$

realvalued

